

**DRUG FACTS**

Active ingredient(s)

Purpose

**Use(s)**

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat
- nasal congestion
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

**Warnings**

**Do not use**

- if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- thyroid disease
- diabetes
- glaucoma
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- liver or kidney disease. Your doctor should determine if you need a different dose.

**Ask a doctor or pharmacist before use if you are**  
**Ask a doctor or pharmacist before use if you are** taking tranquilizers or sedatives.

**When using this product**

- do not use more than directed**
- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- an allergic reaction to this product occurs. Seek medical help right away.
- you get nervous, dizzy, or sleepless
- symptoms do not improve within 7 days or are accompanied by fever

**If pregnant or breast-feeding**

**Keep out of reach of children**  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

<b>Directions</b>	
<ul style="list-style-type: none"><li>do not break or chew tablet; swallow tablet whole</li></ul>	
adults and children 12 years and over	take 1 tablet every 12 hours; do not take more than 2 tablets in 24 hours.
adults 65 years and over	ask a doctor
children under 12 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

**Other information**

- store between 20° to 25°C (68° to 77°F)
- do not use if individual blister unit is open or torn**
- see back panel for lot number and expiration date

**Drug Facts**

**Inactive ingredients**  
colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide